

PEDIATRIC DRUG DOSING DOSE REFERENCE GUIDE - BLS

<u>DRUG</u>	<u>DOSE</u>	<u>MODE</u>	<u>INTERVAL/ RATE</u>	<u>DOSE PREP.</u>
ALBUTEROL	2.5 mg	Nebulizer	X 1	2.5 mg/3 ml
ATROVENT	0.5 mg	Nebulizer	X 1	0.5 mg/3 ml
GLUCOSE, ORAL ¹ (Age 1 month to 4 years)	¼ tube	PO	as indicated	25 gm/tube
GLUCOSE, ORAL (Age 4 to 8 years)	½ tube	PO	as indicated	25 gm/tube
GLUCOSE, ORAL (Age greater than 8 years)	1 tube	PO	as indicated	25 gm/tube
NALOXONE	1 mg: 0-4 years	IN	as indicated	2 mg/2 ml
NALOXONE	2 mg: > 4 years	IN	as indicated	2 mg/2 ml

1 - Oral Glucose is NOT to be used for patients less than 1 month old.

Copyright 2016 Chicago EMS Medical Directors Consortium

Written: 1/09

Reviewed: 3/09; 5/11; 5/16

Revised: 3/09; 5/11; 5/16

MDC Approval 4/7/09; 6/7/11

IDPH Approval: 7/9/09; 9/29/11

Implementation: 1/1/10; 4/1/12