NARROW COMPLEX TACHYCARDIA - ALS
(Pulse >150)



1 – See ECG procedure (ALS Appendix I-15)

2 – If unconscious/unstable, defer vascular access until after cardioversion.

3 - If conscious, consider analgesia prior to cardioversion: Morphine Sulfate 0.1 mg/kg IVP **OR** **Fentanyl 1 mcg/kg IVP**
≤ 65 years of age - max dose 10 mg **≤ 65 years of age - max dose 100 mcg**
> 65 years of age - max dose 5 mg **> 65 years of age - max dose 50 mcg**

4 - Use antecubital if possible. Also, follow rapid IV injection with immediate 10 ml flush of NS.

5 - If history of reactive airway/asthma/COPD, contact Online Medical Control prior to use of adenosine.

Copyright 2019 Chicago EMS Medical Directors Consortium
Written: 1991
Reviewed: 3/92; 10/95; 2/99; 5/04; 3/08; 5/11; 5/14; 9/18
Revised: 3/92; 10/95; 2/99; 5/04; 3/09; 5/11; 5/14; 9/18
MDC Approval: 9/3/92; 3/99; 6/04; 4/7/09; 6/7/11; 5/19/14; 11/8/18
IDPH Approval: 1991; 11/16/92; 2/20/96; Summer 99; 9/04; 7/9/09; 9/29/11; 6/9/14; 12/18/18
Implementation: 2/12/91; 3/1/93; 5/1/96; 8/1/99; 1/1/05; 1/1/10; 4/1/12; 1/18/19